



TITLE:

Completeness of the Generalized Eigenfunctions for relativistic Schrodinger operators I(Spectral and Scattering Theory and Related Topics)

AUTHOR(S):

Wei, Dabi

CITATION:

Wei, Dabi. Completeness of the Generalized Eigenfunctions for relativistic Schrodinger operators I(Spectral and Scattering Theory and Related Topics). 数理解析研究所講究録 2006, 1510: 169-181

ISSUE DATE:

2006-07

URL:

<http://hdl.handle.net/2433/58576>

RIGHT:

Completeness of the Generalized Eigenfunctions for relativistic Schrödinger operators I

Dabi Wei
魏 大比

April 28, 2006

Abstract

Generalized eigenfunctions of the odd-dimensional ($n \geq 3$) relativistic Schrödinger operator $\sqrt{-\Delta} + V(x)$ with $|V(x)| \leq C\langle x \rangle^{-\sigma}$, $\sigma > 1$, are considered. We compute the integral kernels of the boundary values $R^\pm(\lambda) = (\sqrt{-\Delta} - (\lambda \pm i0))^{-1}$, and prove that the generalized eigenfunctions $\varphi^\pm(x, k) := \varphi_0(x, k) - R^\mp(|k|)V\varphi_0(x, k)$ ($\varphi_0(x, k) := e^{i x \cdot k}$) are bounded for $(x, k) \in \mathbb{R}^n \times \{k | a \leq |k| \leq b\}$, where $[a, b] \subset (0, \infty) \setminus \sigma_p(H)$. This fact, together with the completeness of the wave operators, enables us to obtain the eigenfunction expansion for the absolutely continuous spectrum.

Introduction

This paper considers the odd-dimensional ($n \geq 3$) relativistic Schrödinger operator

$$H = H_0 + V(x), \quad H_0 = \sqrt{-\Delta}, \quad x \in \mathbb{R}^n$$

with a short range potential $V(x)$.

Throughout the paper we assume that $V(x)$ is a real-valued measurable function on \mathbb{R}^n satisfying

$$|V(x)| \leq C\langle x \rangle^{-\sigma}, \quad \sigma > 1.$$

When we deal with the boundness and the completeness of the generalized eigenfunctions, σ will be required to satisfy the assumption $\sigma > (n+1)/2$ and n to be an odd integer with $n \geq 3$.

In general, the schrödinger operator is written as $-\Delta + V(x)$, $x \in \mathbb{R}^n$. In [8], the completeness of the generalized eigenfunctions for operator $-\Delta + V(x)$ was proved. However, it was considered by 3-dimensional case. And, in the case of N-body, the completeness was proved in [11, 12]. When the speed of the particles approach light, we have to consider the relativistic case. the schrödinger operator is written by $\sqrt{-\Delta + m^2} + V(x)$, $x \in \mathbb{R}^n$, where m is the mass of the particle. In recent years, there have been

some works on the decay of eigenfunctions associated to the discrete spectra of these operators [4, 20]. On the asymptotic behaviour of the eigenfunctions of the relativistic N-body Schrödinger operator, some works have been done in [21].

But, like a photon, the zero mass particle exists. Then, the relativistic Schrödinger operator is written by $H = \sqrt{-\Delta} + V(x)$, $x \in \mathbb{R}^n$. H is essentially self adjoint on $C_0^\infty(\mathbb{R}^n)$ [27]. And in the paper [28], T. Umeda considered the 3-dimensional case and proved that the generalized eigenfunctions $\varphi^\pm(x, k)$ are bounded for $(x, k) \in \mathbb{R}^3 \times \{k | k \in \mathbb{R}^3, a \leq |k| \leq b\}$, $[a, b] \subset (0, \infty) \setminus \sigma_p(H)$. In the part II, T. Umeda announced that he will deal with the completeness of the generalized eigenfunctions. But, he was too busy to collect his result.

For the purpose of making a comparison, let us briefly recall some results done before. For $z \in \rho(H)$, the resolvents of H and H_0 will be written as

$$R_0(z) = (H_0 - z)^{-1}, \quad R(z) = (H - z)^{-1}.$$

Clearly, for any $\lambda \in (0, \infty) \setminus \sigma_p(H)$ and $s > 1/2$, there exist the limits (see [2, Theorem 4A])

$$\begin{aligned} R_0^\pm(\lambda) &= \lim_{\mu \downarrow 0} R_0(\lambda \pm i\mu) \quad \text{in } \mathbf{B}(L^{2,s}, H^{1,-s}), \\ R^\pm(\lambda) &= \lim_{\mu \downarrow 0} R(\lambda \pm i\mu) \quad \text{in } \mathbf{B}(L^{2,s}, H^{1,-s}). \end{aligned}$$

Following S. Agmon [1], we define two families of generalized eigenfunctions of H by

$$\varphi^\pm(x, k) := \varphi_0(x, k) - R^\mp(|k|)\{V(\cdot)\varphi_0(\cdot, k)\}(x)$$

for k with $|k| \in (0, \infty) \setminus \sigma_p(H)$. In the paper [28, section 8], T. Umeda considered the 3-dimensional case and proved that the generalized eigenfunctions φ^\pm satisfy

$$\varphi^\pm(x, k) = \varphi_0(x, k) - R_0^\mp(|k|)\{V(\cdot)\varphi^\pm(\cdot, k)\}(x)$$

for $(x, k) \in \mathbb{R}^3 \times \{k | k \in \mathbb{R}^3, a \leq |k| \leq b\}$, $[a, b] \subset (0, \infty) \setminus \sigma_p(H)$, which is called modified Leppmann-Schwinger equations. Moreover, he showed that the generalized eigenfunctions $\varphi^\pm(x, k)$ are bounded for $(x, k) \in \mathbb{R}^3 \times \{k | k \in \mathbb{R}^3, a \leq |k| \leq b\}$, $[a, b] \subset (0, \infty) \setminus \sigma_p(H)$, (see T. Umeda [28, section 9]). T. Umeda [29] announced that he will deal with the completeness of the generalized eigenfunctions.

Under the condition of the odd-dimension ($n \geq 3$), the present paper shows that the same equation is valid,

$$\varphi^\pm(x, k) = \varphi_0(x, k) - R_0^\mp(|k|)\{V(\cdot)\varphi^\pm(\cdot, k)\}(x)$$

for $(x, k) \in \mathbb{R}^n \times \{k | a \leq |k| \leq b\}$, $[a, b] \subset (0, \infty) \setminus \sigma_p(H)$, but when $n > 3$, the resolvent is different from the case $n = 3$. The computations show that there exists some polynomials $a_j(z)$, $b_j(z)$, $c_j(z)$ of z in \mathbb{C} such that the integral kernel of the resolvent of $\sqrt{-\Delta}$ is given by

$$R_0(z)u(x) = \int_{\mathbb{R}^n} g_z(x - y)u(y)dy$$

for $z \in \mathbb{C} \setminus [0, \infty)$, where

$$g_z(x) = -\frac{c_n}{2m}|x|^{-2m} + b_{m-1}(z)M_z(x)|x|^{-(m-1)} \\ + \sum_{j=m}^{2m-1} \left(a_j(z) + b_j(z)M_z(x) + c_j(z)N_z(x) \right) |x|^{-j},$$

and

$$M_z(x) = \frac{1}{|x|} \left\{ \text{ci}(-|x|z) \sin(|x|z) - \text{si}(-|x|z) \cos(|x|z) \right\}, \\ N_z(x) = \left\{ \text{ci}(-|x|z) \cos(|x|z) + \text{si}(-|x|z) \sin(|x|z) \right\}.$$

For the definitions of the cosine and sine integral functions $\text{ci}(z)$ and $\text{si}(z)$, see section 5. We compute the limit $g_\lambda^\pm(x) := \lim_{\mu \downarrow 0} g_{\lambda \pm i\mu}(x)$ as follows,

$$g_\lambda^\pm(x) = \left\{ a_{2m}(\lambda) + b_{2m}(e^{\pm i\lambda|x|} + m_\lambda(x)) \right\} |x|^{-2m} \\ + \sum_{j=m}^{2m-1} a_j(\lambda) |x|^{-j} + \sum_{j=m}^{2m-1} b_j(\lambda) (e^{\pm i\lambda|x|} + m_\lambda(x)) |x|^{-j} \\ + \sum_{j=m}^{2m-1} c_j(\lambda) (e^{\pm i(\lambda|x| + \pi/2)} + n_\lambda(x)) |x|^{-j},$$

where

$$m_\lambda(x) = \text{ci}(\lambda|x|) \sin(\lambda|x|) + \text{si}(\lambda|x|) \cos(\lambda|x|), \\ n_\lambda(x) = \text{ci}(\lambda|x|) \cos(\lambda|x|) - \text{si}(\lambda|x|) \sin(\lambda|x|).$$

We then prove that the generalized eigenfunctions $\varphi^\pm(x, k)$ are bounded for $(x, k) \in \mathbb{R}^n \times \{k \mid a \leq |k| \leq b\}$, $[a, b] \subset (0, \infty) \setminus \sigma_p(H)$, with

$$R_0^\pm(\lambda)u(x) = \int_{\mathbb{R}^n} g_\lambda^\pm(x-y)u(y)dy.$$

$H = H_0 + V$ defines a selfadjoint operator in $L^2(\mathbb{R}^n)$, whose domain is $H^1(\mathbb{R}^n)$ (see section 2, or T. Umeda [27, Theorem 5.8]). Moreover, H is essentially selfadjoint on $C_0^\infty(\mathbb{R}^n)$ (see T. Umeda [27]). It follows from Reed-Simon [22, P113, Corollary 2] that

$$\sigma_e(H) = \sigma_e(H_0) = [0, \infty).$$

The fact that $\sigma_p(H) \cap (0, \infty)$ is a discrete set was first proved by B. Simon [23, Theorem 2.1]. He also proved that each eigenvalue in the set $\sigma_p(H) \cap (0, \infty)$ has finite multiplicity [23, Theorem 2.1]. From V. Enss's idea (see V. Enss [5]), we obtain that the wave operators W_\pm defined by

$$W_\pm = \lim_{t \rightarrow \infty} e^{itH} e^{-itH_0}$$

are complete. Finally, by the idea of H. Kitada [12] and S.T. Kuroda [15], we obtain the completeness of the generalized eigenfunctions as follows.

Theorem Assume the dimension n ($n \geq 3$) is an odd integer, $\sigma > (n+1)/2$, $s > n/2$ and $[a, b] \subset (0, \infty) \setminus \sigma_p(H)$. For $u \in L^{2,s}(\mathbb{R}^n)$, let \mathcal{F}_\pm be defined by

$$\mathcal{F}_\pm u(k) := (2\pi)^{-n/2} \int_{\mathbb{R}^n} u(x) \overline{\varphi^\pm(x, k)} dx.$$

Then for an arbitrary $L^{2,s}(\mathbb{R}^n)$ -function $f(x)$,

$$E_H([a, b])f(x) = (2\pi)^{-n/2} \int_{a \leq |k| \leq b} \mathcal{F}_\pm f(k) \varphi^\pm(x, k) dk$$

where E_H is the spectral measure on H .

The plan of the paper In section 1, we construct generalized eigenfunctions of $\sqrt{-\Delta} + V(x)$ on \mathbb{R}^n . We compute the resolvent kernel of $\sqrt{-\Delta}$ on \mathbb{R}^n in the integral form in section 2. Section 3 prove the generalized eigenfunctions are bounded in the case of odd-dimension $n \geq 3$. We studies the asymptotic completeness of wave operators in section 4. In the last section 5, we deal with the completeness of the generalized eigenfunctions. We explained about the theorems without proving in this paper for the limitation of the number of pages.

Notation We introduce the notation which will be used in the present paper.

For $x \in \mathbb{R}^n$, $|x|$ denotes the Euclidean norm of x and $\langle x \rangle = \sqrt{1 + |x|^2}$. The Fourier transform of a function u is denoted by $\mathcal{F}u$ or \hat{u} , and is defined by

$$\mathcal{F}u(\xi) = \hat{u}(\xi) = (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{-ix \cdot \xi} u(x) dx.$$

For s and l in \mathcal{R} , we define the weighted L^2 -space and the weighted Sobolev space by

$$L^{2,s}(\mathbb{R}^n) = \{f | \langle x \rangle^s f \in L^2(\mathbb{R}^n)\}, \quad H^{l,s}(\mathbb{R}^n) = \{f | \langle x \rangle^s \langle D \rangle^l f \in L^2(\mathbb{R}^n)\}$$

respectively, where D stands for $-i\partial/\partial x$ and $\langle D \rangle = \sqrt{1 + |D|^2} = \sqrt{1 - \Delta}$. The inner products and the norm in $L^{2,s}(\mathbb{R}^n)$ and $H^{l,s}(\mathbb{R}^n)$ are given by

$$(f, g)_{L^{2,s}} = \int_{\mathbb{R}^n} \langle x \rangle^{2s} f(x) \overline{g(x)} dx, \quad (f, g)_{H^{l,s}} = \int_{\mathbb{R}^n} \langle x \rangle^{2s} \langle D \rangle^l f(x) \overline{\langle D \rangle^l g(x)} dx,$$

$$\|f\|_{L^{2,s}} = \{(f, f)_{L^{2,s}}\}^{1/2}, \quad \|f\|_{H^{l,s}} = \{(f, f)_{H^{l,s}}\}^{1/2},$$

respectively. For $s = 0$ we write

$$(f, g) = (f, g)_{L^{2,0}} = \int_{\mathbb{R}^n} f(x) \overline{g(x)} dx, \quad \|f\|_{L^2} = \|f\|_{L^{2,0}}.$$

For a pair of $f \in L^{2,-s}(\mathbb{R}^n)$ and $g \in L^{2,s}(\mathbb{R}^n)$, we also define $(f, g) = \int_{\mathbb{R}^n} f(x) \overline{g(x)} dx$.

By $C_0^\infty(\mathbb{R}^n)$ we mean the space of C^∞ -functions of compact support. By $\mathcal{S}(\mathbb{R}^n)$ we mean the Schwartz space of rapidly decreasing functions, and by $\mathcal{S}'(\mathbb{R}^n)$ the space of tempered distributions.

The operator $\sqrt{-\Delta}e^{ix \cdot k}$ is formally defined by

$$\int_{\mathbb{R}^n} e^{ix \cdot \xi} |\xi| \delta(\xi - k) d\xi,$$

where $\delta(x)$ is the Dirac's delta function. As the symbol $|\xi|$ of $\sqrt{-\Delta}$ is singular at the origin $\xi = 0$, making sense of the expression $\sqrt{-\Delta}e^{ix \cdot k}$ is one of the main tasks in the present paper.

For a pair of Hilbert spaces \mathcal{H} and \mathcal{K} , $\mathbf{B}(\mathcal{H}, \mathcal{K})$ denotes the Banach space of all bounded linear operators from \mathcal{H} to \mathcal{K} .

For a selfadjoint operator H in a Hilbert space, $\sigma(H)$ and $\rho(H)$ denote the spectrum of H and the resolvent set of H , respectively. The essential spectrum, the continuous spectrum and the absolutely continuous spectrum of H will be denoted by $\sigma_e(H)$, $\sigma_c(H)$, and $\sigma_{ac}(H)$ respectively. E_H denotes the spectral measure on T , and $E_H(\lambda) = E_H((-\infty, \lambda])$, $E_H((a, b]) = E_H(b) - E_H(a)$.

The continuous subspace and the absolutely continuous subspace of H will be denoted by \mathcal{H}_c , \mathcal{H}_{ac} , respectively.

1 Generalized eigenfunction

We construct in this section generalized eigenfunctions of $\sqrt{-\Delta} + V(x)$ on \mathbb{R}^n , and show that the generalized eigenfunctions satisfy the equation

$$\varphi^\pm(x, k) = \varphi_0(x, k) - R_0^\mp(|k|)V\varphi^\pm(x, k),$$

where $R_0(z)$ is the resolvent of $H_0 = \sqrt{-\Delta}$ defined by

$$R_0(z) := (H_0 - z)^{-1} = \mathcal{F}^{-1}(|\xi| - z)^{-1}\mathcal{F},$$

and $\varphi_0(x, k)$ is defined by

$$\varphi_0(x, k) = e^{ix \cdot k}.$$

Similarly $R(z)$ is the resolvent of $H = \sqrt{-\Delta} + V(x)$ on \mathbb{R}^n and we assume that $V(x)$ is a real-valued measurable function on \mathbb{R}^n satisfying $|V(x)| < C\langle x \rangle^{-\sigma}$ for some $\sigma > 1$. To show the above equation for eigenfunctions, we use two theorems demonstrated by Ben-Artzi and Nemirovki. (see [2, Section 2 and Theorem 4A])

Theorem 1.1 (Ben-Artzi and Nemirovki) Let $s > 1/2$. Then

- (1) For any $\lambda > 0$, there exist the limits $R_0^\pm(\lambda) = \lim_{\mu \downarrow 0} R_0(\lambda \pm i\mu)$ in $\mathbf{B}(L^{2,s}, H^{1,-s})$.
- (2) The operator-valued functions $R_0^\pm(z)$ defined by

$$R_0^\pm(z) = \begin{cases} R_0(z) & \text{if } z \in \mathbb{C}^\pm \\ R_0^\pm(\lambda) & \text{if } z = \lambda > 0 \end{cases}$$

are $B(L^{2,s}, H^{1,-s})$ -valued continuous functions, where \mathbb{C}^+ and \mathbb{C}^- are the upper and the lower half-planes respectively: $\mathbb{C}^\pm = \{z \in \mathbb{C} \mid \pm \operatorname{Im} z > 0\}$.

Theorem 1.2 (Ben-Artzi and Nemirovki) Let $s > 1/2$ and $\sigma > 1$. Then

(1) The continuous spectrum $\sigma_c(H) = [0, \infty)$ is absolutely continuous, except possibly for a discrete set of embedded eigenvalues $\sigma_p(H) \cap (0, \infty)$, which can accumulate only at 0 and ∞ .

(2) For any $\lambda \in (0, \infty) \setminus \sigma_p(H)$, there exist the limits

$$R^\pm(\lambda) = \lim_{\mu \downarrow 0} R(\lambda \pm i\mu) \quad \text{in} \quad B(L^{2,s}, H^{1,-s}).$$

(3) The operator-valued functions $R^\pm(z)$ defined by

$$R^\pm(z) = \begin{cases} R(z) & \text{if } z \in \mathbb{C}^\pm \\ R^\pm(\lambda) & \text{if } z = \lambda > 0 \setminus \sigma_p(H) \end{cases}$$

are $B(L^{2,s}, H^{1,-s})$ -valued continuous functions.

The main results of this section are

Theorem 1.3 Let $\sigma > (n+1)/2$, if $|k| \in (0, \infty) \setminus \sigma_p(H)$, then generalized eigenfunctions

$$\varphi^\pm(x, k) := \varphi_0(x, k) - R^\mp(|k|)\{V(\cdot)\varphi_0(\cdot, k)\}(x)$$

satisfy the equation

$$(\sqrt{-\Delta_x} + V(x))u = |k|u \quad \text{in} \quad \mathcal{S}'(\mathbb{R}_x^n)$$

where $\varphi_0(x, k)$ is defined by $\varphi_0(x, k) = e^{ix \cdot k}$.

Theorem 1.4 Let $\sigma > (n+1)/2$. If $|k| \in (0, \infty) \setminus \sigma_p(H)$, and $n/2 < s < \sigma - 1/2$, then we have

$$\varphi^\pm(x, k) = \varphi_0(x, k) - R_0^\mp(|k|)\{V(\cdot)\varphi^\pm(\cdot, k)\}(x) \quad \text{in} \quad L^{2,-s}(\mathbb{R}^n).$$

2 The integral kernel of the resolvents of H_0

This section is devoted to computing the resolvent kernel of $H_0 = \sqrt{-\Delta}$ on \mathbb{R}^n , where $n = 2m + 1$, $m \geq 1$ and $m \in \mathbb{N}$. Then we compute the limit of $g_z(x)$ as $\mu \downarrow 0$, where $z = \lambda + i\mu$ and $\lambda > 0$, and study the properties of the integral operator G_λ^\pm . In this section we suppose that (cf. [6, p. 269, Formula (46) and (47)])

(1) $n = 2m + 1$, $m \geq 1$ and $m \in \mathbb{N}$,

$$(2) \quad M_z(x) = \int_0^\infty e^{tz} \frac{1}{t^2 + |x|^2} dt = \frac{1}{|x|} \left\{ \operatorname{ci}(-|x|z) \sin(|x|z) - \operatorname{si}(-|x|z) \cos(|x|z) \right\},$$

$$N_z(x) = \int_0^\infty e^{tz} \frac{t}{t^2 + |x|^2} dt = \operatorname{ci}(-|x|z) \cos(|x|z) + \operatorname{si}(-|x|z) \sin(|x|z),$$

(3) $m_\lambda(x) = \operatorname{ci}(\lambda|x|) \sin(\lambda|x|) + \operatorname{si}(\lambda|x|) \cos(\lambda|x|),$

$$n_\lambda(x) = \operatorname{ci}(\lambda|x|) \cos(\lambda|x|) - \operatorname{si}(\lambda|x|) \sin(\lambda|x|).$$

Where $\text{ci}(x)$ and $\text{si}(x)$ are defined by

$$\text{ci}(x) = \int_x^\infty \frac{\cos t}{t} dt, \quad \text{si}(x) = - \int_x^\infty \frac{\sin t}{t} dt, \quad x > 0.$$

We see that $\text{si}(x)$ has an analytic continuation $\text{si}(z)$ (see [6, P145]),

$$\text{si}(z) = -\frac{\pi}{2} + \sum_{m=0}^{\infty} \frac{(-1)^m}{(2m+1)!(2m+1)} z^{2m+1} \quad (2.1)$$

The cosine integral function $\text{ci}(x)$ has an analytic continuation $\text{ci}(z)$, which is a many-valued function with a logarithmic branch-point at $z = 0$ (see [6, P145]). In this paper, we choose the principal branch

$$\text{ci}(z) = -\gamma - \text{Log} z - \sum_{m=1}^{\infty} \frac{(-1)^m}{(2m)!2m} z^{2m}, \quad z \in \mathbb{C} \setminus (-\infty, 0], \quad (2.2)$$

where γ is the Euler's constant. The main theorems are

Theorem 2.1 Let $n \geq 3$, $\text{Re} z < 0$, then

$$R_0(z)u = G_z u$$

for all $u \in C_0^\infty(\mathbb{R}^n)$, where

$$\begin{aligned} G_z u(x) &= \int_{\mathbb{R}^n} g_z(x-y)u(y)dy, \quad g_z(x) = \int_0^\infty e^{tz} \frac{c_n t}{(t^2 + |x|^2)^{\frac{n+1}{2}}} dt, \\ c_n &= \pi^{-\frac{n+1}{2}} \Gamma\left(\frac{n+1}{2}\right), \quad \Gamma(x) = \int_0^\infty s^{x-1} e^{-s} ds. \end{aligned} \quad (2.3)$$

Theorem 2.2 Let $n = 2m + 1$, $m \geq 1$ ($m \in \mathbb{N}$) and $s > 1/2$, $u \in L^{2,s}(\mathbb{R}^n)$. Let $[a, b] \subset (0, \infty)$ and $\lambda \in [a, b]$.

(1) There exist some functions $a_j(\lambda)$, $b_j(\lambda)$, $c_j(\lambda)$ which are polynomials of λ for $j = m, m+1, \dots, 2m$,

$$\begin{aligned} R_0^\pm(\lambda)u(x) &= G_\lambda^\pm u(x) = \int_{\mathbb{R}^n} g_\lambda^\pm(x-y)u(y)dy \\ g_\lambda^\pm(x) &:= \lim_{\mu \downarrow 0} g_{\lambda \pm i\mu}(x) = \{a_{2m}(\lambda) + b_{2m}(e^{\pm i\lambda|x|} + m_\lambda(x))\} |x|^{-2m} \\ &\quad + \sum_{j=m}^{2m-1} a_j(\lambda) |x|^{-j} + \sum_{j=m}^{2m-1} b_j(\lambda) (e^{\pm i\lambda|x|} + m_\lambda(x)) |x|^{-j} \\ &\quad + \sum_{j=m}^{2m-1} c_j(\lambda) (e^{\pm i(\lambda|x| + \pi/2)} + n_\lambda(x)) |x|^{-j}, \end{aligned}$$

where $R_0^\pm(\lambda) := \lim_{\mu \downarrow 0} R_0(\lambda \pm i\mu)$.

(2) There exist some positive constants C_{abj} for $j = m, m+1, \dots, 2m$ such that

$$|R_0^\pm(\lambda)u(x)| = |G_\lambda^\pm u(x)| \leq \sum_{j=m}^{2m} |D_j u(x)|,$$

$$D_j(\lambda)u(x) := C_{abj} \int_{\mathbb{R}^n} |x-y|^{-j} u(y) dy.$$

3 Boundness of the generalized eigenfunctions

In this section, we assume that n , $V(x)$ and k satisfy the next inequalitys:

- (1) $n = 2m + 1$ ($m \in \mathbb{N}$) and $m \geq 1$
- (2) $|V(x)| \leq C\langle x \rangle^{-\sigma}$, $\sigma > \frac{n+1}{2}$
- (3) $k \in \left\{k \mid a \leq |k| \leq b\right\}$ and $[a, b] \subset (0, \infty) \setminus \sigma_p(H)$.

Applying Theorem 1.4, we see that generalized eigenfunction $\varphi^\pm(x, k)$ defined by

$$\varphi^\pm(x, k) = \varphi_0(x, y) - R^\mp(|k|)\{V(\cdot)\varphi_0(\cdot, k)\}(x), \quad (3.1)$$

satisfies the equation

$$\varphi^\pm(x, k) = \varphi_0(x, k) - R_0^\mp(|k|)\{V(\cdot)\varphi^\pm(\cdot, k)\}(x), \quad (3.2)$$

where $\varphi_0(x, k) = e^{ix \cdot k}$.

In this section, we let $\{D_j V(\cdot)\varphi^\pm(\cdot, k)\}(x)$ be denoted by $D_j V(x)\varphi^\pm(x, k)$. Moreover, let $V(x)D_{j_r} V(x)D_{j_{r-1}} \cdots V(x)D_{j_1} V(x)\varphi^\pm(x, k)$, be denoted by

$$\left(\prod_{p=1}^r V(x)D_{j_p} \right) \{V(x)\varphi^\pm(x, k)\}.$$

This section proves boundness of the generalized eigenfunctions. And the main theorem is

Theorem 3.1 Let $n = 2m + 1$, $m \geq 1$ ($m, n \in \mathbb{N}$), and $[a, b] \subset (0, \infty) \setminus \sigma_p(H)$. Then there exists a constant C_{ab} such that generalized eigenfunctions defined by $\varphi^\pm(x, k) := \varphi_0(x, y) - R^\mp(|k|)\{V(\cdot)\varphi_0(\cdot, k)\}(x)$ satisfy

$$|\varphi^\pm(x, k)| \leq C_{ab},$$

for all $(x, k) \in \mathbb{R}^n \times \{a \leq |k| \leq b\}$, where $\varphi_0(x, k) = e^{ix \cdot k}$.

To prove the main theorem, we gave the lemmas as follows.

Lemma 3.1 Let $m+1 \leq j \leq 2m$ ($j \in \mathbb{N}$) and $p > n/(n-j)$. If $u(x, k) \in L^2(\mathbb{R}_x^n) \cap L^p(\mathbb{R}_x^n)$, and $\|u(x, k)\|_{L^2} \leq C'_{ab}$, $\|u(x, k)\|_{L^p} \leq C''_{ab}$, (C'_{ab} and C''_{ab} are positive

constants) for all $(x, k) \in \mathbb{R}^n \times \{a \leq |k| \leq b\}$, then there exists a positive constant C_{ab} , such that

$$|D_j u(x, k)| \leq C_{ab},$$

for all $(x, k) \in \mathbb{R}^n \times \{a \leq |k| \leq b\}$.

Lemma 3.2 Let $r, j_p \in \mathbb{N}$ and $s > 1/2$. If $m+1 \leq j_p \leq 2m$ for $1 \leq p \leq r$, then

$$\left(\prod_{p=1}^r V(x) D_{j_p} \right) \{V(x) \varphi^\pm(x, k)\} \in L^{2,s}(\mathbb{R}_x^n)$$

for all $r \in \mathbb{N}$. Moreover, there exists a positive constant C_{ab} , such that

$$\left\| \left(\prod_{p=1}^r V(x) D_{j_p} \right) \{V(x) \varphi^\pm(x, k)\} \right\|_{L^{2,s}} \leq C_{ab}$$

for all $(x, k) \in \mathbb{R}^n \times \{a \leq |k| \leq b\}$.

Lemma 3.3 Let $0 < \alpha < n$, $1 < p < q < \infty$ and $f \in L^p(\mathbb{R}^n)$. Let $I_\alpha f(x)$ be defined by $I_\alpha f(x) := \int_{\mathbb{R}^n} |x-y|^{-n+\alpha} f(y) dy$. If $1/q = 1/p - \alpha/n$, there exists a positive constant C_{pq} , such that

$$\|I_\alpha f\|_{L^q} \leq C_{pq} \|f\|_{L^p}.$$

For the proof of the theorem, see [24, P119].

Lemma 3.4 Let $r \in \mathbb{N}$. If $m+1 \leq j_p \leq 2m$ ($1 \leq q \leq r$), and $2 \sum_{p=1}^q j_p > (2q-1)n$

for all $q \leq r$, then

$$\left(\prod_{p=1}^r V(x) D_{j_p} \right) \{V(x) \varphi^\pm(x, k)\} \in L^{\frac{2n}{2 \sum_{p=1}^r j_p - (2r-1)n}}(\mathbb{R}_x^n)$$

for all $r \leq n-1$. Moreover, there exists a positive constant C_{ab} , such that

$$\left\| \left(\prod_{p=1}^r V(x) D_{j_p} \right) \{V(x) \varphi^\pm(x, k)\} \right\|_{L^{\frac{2n}{2 \sum_{p=1}^r j_p - (2r-1)n}}} \leq C_{ab}$$

for all $(x, k) \in \mathbb{R}^n \times \{a \leq |k| \leq b\}$.

Lemma 3.5 Let $r \in \mathbb{N}$ and $r \leq n$. If $m \leq j_p \leq 2m$ for all $1 \leq p \leq r$, then there exist a positive constant C_{ab} , such that

$$\sum_{2 \sum_{p=1}^r j_p < (2r-1)n} \left| D_{j_r} \left(\prod_{p=1}^{r-1} V(x) D_{j_p} \right) \{V(x) \varphi^\pm(x, k)\} \right| \leq C_{ab},$$

for all $(x, k) \in \mathbb{R}^n \times \{a \leq |k| \leq b\}$.

4 Asymptotic completeness

We investigate the asymptotic completeness of wave operators in this section. We assume that the potential $V(x)$ is a real-valued measurable function on \mathbb{R}^n satisfying

$$|V(x)| \leq C\langle x \rangle^{-\sigma} \quad \sigma > 1 \quad (4.1)$$

Under this assumption, it is obvious that V is a bounded selfadjoint operator in $L^2(\mathbb{R}^n)$, and that $H = H_0 + V$ defines a selfadjoint operator in $L^2(\mathbb{R}^n)$, whose domain is $H^1(\mathbb{R}^n)$ (see T. Umeda [27, Theorem 5.8]). Moreover H is essentially selfadjoint on $C_0^\infty(\mathbb{R}^n)$ (see T. Umeda [27]). Since V is relatively compact with respect to H_0 , it follows from Reed-Simon [22, P113, Corollary 2] that

$$\sigma_e(H) = \sigma_e(H_0) = [0, \infty).$$

In this section, we prove the next main theorem with V. Enss's idea (see V. Enss [5] and H. Isozaki [9]).

Theorem 4.1 Let $H_0 = \sqrt{-\Delta}$, $H = H_0 + V(x)$ and $V(x)$ satisfying (4.1). Then there exists the limits

$$W_\pm = \lim_{t \rightarrow \pm\infty} e^{itH} e^{-itH_0},$$

and the asymptotic completeness hold:

$$\mathcal{R}(W_\pm) = \mathcal{H}_{ac}(H).$$

5 Eigenfunction expansions

In this section, we assume that the dimension n is an odd integer, $n \geq 3$, and $\sigma > (n+1)/2$. We consider the completeness of the generalized eigenfunction in this section. The main idea is the same as the idea in H. Kitada [12] and S.T. Kuroda [15], besides, in this section, we use the method in T. Ikebe [8, section 11]. It is known that

$$\sigma_e(H) = \sigma_e(H_0) = [0, \infty).$$

We need to remark that $\sigma_p(H) \cap (0, \infty)$ is a discrete set. This fact was first proved by B. Simon [23, Theorem 2.1]. Moreover, B. Simon [23, Theorem 2.1] proved that each eigenvalue in the set $\sigma_p(H) \cap (0, \infty)$ has finite multiplicity.

The main theorem is

Theorem 5.1 Assume the dimension n ($n \geq 3$) is an odd integer, $\sigma > (n+1)/2$, $s > n/2$ and $[a, b] \subset (0, \infty) \setminus \sigma_p(H)$. For $u \in L^{2,s}(\mathbb{R}^n)$, let \mathcal{F}_\pm be defined by

$$\mathcal{F}_\pm u(k) := (2\pi)^{-n/2} \int_{\mathbb{R}^n} u(x) \overline{\varphi^\pm(x, k)} dx. \quad (5.1)$$

For an arbitrary $L^{2,s}(\mathbb{R}^n)$ -function $f(x)$,

$$E_H([a, b])f(x) = (2\pi)^{-n/2} \int_{a \leq |k| \leq b} \mathcal{F}_\pm f(k) \varphi^\pm(x, k) dk,$$

where E_H is the spectral measure on H , and $\varphi^\pm(x, k)$ are defined in Theorem 1.3.

Acknowledgements The author wishes to express his sincere thanks to Professor H. Kitada for his encouraging and stimulating discussions with him, and thanks to Professor T. Umeda for his encouraging communications. The author also wishes to express his sincere thanks to his family and Ms. T. Cai for their constant supports.

References

- [1] S. Agmon, *Spectral properties of Schrödinger operators and scattering theory*, Ann. Scuola Norm. Sup. Pisa, **4-2** (1975), 151-218.
- [2] M. Ben-Artzi and J. Nemirovski, *Remarks on relativistic Schrödinger operators and their extensions*, Ann. Inst. Henri Poincaré, Phys. théor. **67** (1997), 29-39.
- [3] M. Ben-Artzi, *Regularity and Smoothing for some Equations of Evolution in Non-linear Partial Differential Equations and Their Applications*, Vol. **11**, (1994), 1-12.
- [4] R. Carmona, W.C. Masters and B. Simon, *Relativistic Schrödinger operators: asymptotic behavior of the eigenfunctions*, J. Funct. Analysis **91** (1990), 117-142.
- [5] V. Enss, *Asymptotic completeness for quantum-mechanical potential scattering I, Short range potentials*, Commun. Math. Phys. **61**, (1978), 258-291.
- [6] A. Erdélyi ed., *Tables of Integral Transforms vol. 1*, McGraw-Hill (1952).
- [7] T. Ikebe and Y. Saito, *Limiting absorption method and absolute continuity for the Schrödinger operators*, J. Math. Kyoto Univ. **7** (1972), 513-542.
- [8] T. Ikebe, *Eigenfunction expansions associated with the Schrödinger operators and their applications to scattering theory*, Arch. Rational Mech. Anal., **5** (1960), 1-34.
- [9] H. Isozaki, *Many-body Schrödinger equation*, Springer Tokyo (2004). (In Japanese).
- [10] H. Kitada, *Quantum Mechanics*, <http://kims.ms.u-tokyo.ac.jp/quantum.html> (2003).
- [11] H. Kitada, *Fundamental solutions and eigenfunction expansions for Schrödinger Operators I: Fundamental Solutions*, Math. Z. **198**, (1988), 181-190.
- [12] A. Jensen and H. Kitada, *Fundamental solutions and eigenfunction expansions for Schrödinger Operators II: Eigenfunction Expansions*, Math. Z. **199**, (1988), 1-13.
- [13] H. Kitada and H. Kumano-go, *A family of Fourier integral operators and the fundamental solution for a Schrödinger equation*, Osaka J. Math. **18**, (1981), 291-360.
- [14] H. Kumano-go, *Pseudo-differential Operators*, MIT Press (1981).

- [15] S.T. Kuroda, *Spectral theory II*, Iwanami Shoten, Tokyo, (1979). (In Japanese).
- [16] S.T. Kuroda, *Theory of simple scattering and eigenfunction expansions. In: functional analysis and related fields*, Berlin-Heidelberg-New York: Springer (1970), 99-131.
- [17] S.T. Kuroda, *Scattering theory for differential operators I*, J. Math. Soc. Japan **25** (1973), 75-104.
- [18] S.T. Kuroda, *Scattering theory for differential operators II*, J. Math. Soc. Japan **25** (1973), 222-234.
- [19] M.Nagase and T.Umeda, *On the essential self-adjointness of pseudo-differential operators*, Proc. Japan Acad. 64 Ser. A (1988), 94-97.
- [20] F. Nardini, *Exponential decay for the eigenfunctions of the two body relativistic Hamiltonians*, J. D 'Analyse Math. 47 (1986), 87.109.
- [21] F. Nardini, *On the asymptotic behaviour of the eigenfunctions of the relativistic N-body Schrödinger operator*, Boll. Un. Mat. Ital. A (7) 2 (1988), 365.369.
- [22] M. Reed and B. Simon, *Methods of Modern Mathematical Physics IV: Analysis of Operators*, Academic Press (1978).
- [23] B. Simon, *Phase space analysis of simple scattering systems: Extensions of some work of Enss*, Duke Math. J., **46** (1979), 119-168.
- [24] E.M. Stein, *Singular integrals and differentiability properties of functions*, Princeton University Press (1970).
- [25] R. Strichartz, *A Guide to Distribution Theory and Fourier Transforms*, World Scientific (2003).
- [26] T.Umeda, *Radiation conditions and resolvent estimates for relativistic Schrödinger*, Ann. Inst. Henri Poincare, Phys. theor. 63(1995), 277-296.
- [27] T.Umeda, *The action of $\sqrt{-\Delta}$ on weighted Sobolev spaces*, Lett. Math. Phys. 54 (2000), 301-313.
- [28] T. Umeda, *Generalized eigenfunctions of relativistic Schrödinger operators I*, Department of Mathematics Himeji Institute of Technology, Japan. (2001).
- [29] T. Umeda, *Generalized eigenfunctions of relativistic Schrödinger operators II*, private communication.
- [30] T.Umeda, *Eigenfunction expansions associated with relativistic Schrödinger operators*, in Partial Differential Equations and Spectral Theory, eds. M. Demuth and B. W. Schulze, Operator Theory: Advances and Applications **126** (2001), 315-319.

Dabi Wei : Department of Information Physics and Computing, Graduate School of Information Science and Technology, University of Tokyo Hongou 7-3-1, bunkyo-ku, Tokyo 113-8656, Japan. *Email* : dabi@alab.t.u-tokyo.ac.jp